

The Pediatric Infectious Disease Journal Publish Ahead of Print

DOI: 10.1097/INF.0000000000001967

Quantifying the Impact of Maternal Influenza Vaccination: Beyond Laboratory-Confirmed Efficacy

Kathleen M. Neuzil, MD, MPH

University of Maryland School of Medicine Center for Vaccine Development

Corresponding Author: Kathleen M. Neuzil, MD, MPH, Professor of Medicine and Pediatrics, Director, Center for Vaccine Development, University of Maryland School of Medicine, 685 West Baltimore St. HSF1-Room 480, Baltimore, MD 21201, 410-706-5328, kneuzil@som.umaryland.edu;kneuzil@medicine.umaryland.edu

Abbreviated title/Running head: Quantifying the Impact of Maternal Influenza Vaccination

In many high-income countries, pregnant women have long been prioritized to receive influenza vaccine, based on expected benefits to the woman and the infant. In a global evaluation based on data collected in 2014, the World Health Organization (WHO) reported that 75% of high income countries had influenza vaccine policies that targeted pregnant women. Unfortunately, most low and middle-income countries do not use influenza vaccine. In that same WHO survey, of 49 low and middle-income countries eligible for Gavi support, only 4 (8%) had any influenza vaccine policy and only 2 (4%) targeted pregnant women.¹

A number of factors influence vaccine policy and uptake, including demonstrated disease burden and vaccine impact, financing mechanisms, and logistical considerations. Particularly for low income countries, with competing health priorities, demonstration of impact of the vaccine on severe disease is critical. Unfortunately, few definitive data exist for the vaccine-preventable burden of severe influenza illness in low-resource settings. Demonstrating such burden for influenza can be particularly challenging, given that influenza has substantial variability from season-to-season, and that most prospective studies are not conducted over multiple seasons or sufficiently powered for less common, more severe outcomes. Further, influenza diagnostics are not routinely used in low resource settings. Even when diagnostics are employed, severe respiratory disease may occur as a consequence of earlier influenza infection and thus influenza virus may not be detectable at the time the severe illness occurs.²

Vaccine probe studies are an alternative way to establish disease burden.^{2,3} The premise is that differences in outcomes identified in a randomized, controlled vaccine trial can be assumed to be due to the disease targeted by the vaccine, even in the absence of a laboratory confirmation of the outcome. Probe studies have been instrumental in establishing the burden of all-cause pneumonia and deaths due to *Streptococcus pneumoniae* and *Haemophilus influenzae* type b, and

the burden of severe gastroenteritis attributable to rotavirus.^{3,4} Such studies have influenced policy and financing decisions for those vaccines.

From 2011-2014, three randomized trials of maternal influenza vaccine were conducted in South Africa, Mali and Nepal. These landmark studies represented a major investment to understand the potential benefits of maternal influenza vaccine in preventing maternal and infant disease. While the pre-stated primary and secondary outcomes differed among the trials, in general the trials were powered to detect laboratory-confirmed influenza of any severity. Vaccine was well-tolerated in all 3 studies, and all studies reported significant efficacy against laboratory-confirmed disease in mothers and infants, although results differed by setting and season.^{5, 6, 7}

A study in this issue of the *Pediatric Infectious Disease Journal* reports on a pooled analysis of those 3 randomized controlled trials to determine the effect of maternal influenza vaccine on severe clinical pneumonia of any etiology in the infants.⁸ Severe pneumonia was defined as hospitalization for any lower respiratory disease in South Africa and by Integrated Management of Childhood Illness (ICMI) criteria in Nepal and Mali. Overall, the authors report an adjusted incidence of 87.7 and 70.7 cases of severe pneumonia per 1000 infant-years, respectively, in maternal control and influenza vaccine groups for an overall efficacy of 31.3%. Efficacy estimates were greater during times of influenza circulation as compared to no circulation. In analyses by site, vaccine efficacy estimates in Nepal and South Africa were similar to the overall pooled estimate, while no effect of maternal influenza vaccination on infant pneumonia was demonstrated in the Mali site.

The inherent variability of influenza complicates execution and interpretation of clinical vaccine trials. This is true for this pooled evaluation as well. During the period of the trial, the sites had variability in influenza circulation (different circulating strains, different timing of peak

season in relation to when vaccine was given); population (gestational age at receipt of vaccine, sociodemographic characteristics), and vaccine used (different formulations, different degree of “match” and different comparator vaccines). The trial size and multi-year design were strengths, but do not completely overcome these limitations. In addition, other differences in study design, including the definition of severe pneumonia, add to the heterogeneity of the data and cloud interpretation of the results. Unadjusted incidence rates are not provided and would be helpful in assessing the generalizability to other settings and comparability to other studies. In spite of these challenges, these large randomized controlled trials provide a rich dataset that should be further explored. For example, Tapia and colleagues in Mali demonstrated that maternal influenza immunization provided protection that corresponded with infant age and level of maternal antibody. Protection was highest in the first 2 months of life. Thereafter, as influenza antibody titers diminished, efficacy decreased, until it was no longer evident at month 6 of follow-up.⁶ One would expect any vaccine protection against severe pneumonia to follow a similar pattern – with potentially a delay for pneumonia if the upper respiratory infection leads to subsequent lower respiratory viral or bacterial disease. Either way, looking at the magnitude of the effect by age of the infant would enhance the biological plausibility for this finding.

Nunes and colleagues did evaluate the effect of maternal vaccination on infant lower respiratory tract disease in a single site analysis from the South Africa trial.⁹ In that study, the incidence of hospitalizations for acute lower respiratory infection was lower in infants born to influenza vaccine recipients (3.4 per 1000 infant-months) compared with placebo recipients (6.0 per 1000 infant-months) with a vaccine efficacy of 43.1% ($P = .050$). The full effect of maternal vaccination was seen in the first 90 days of life; the incidence of acute lower respiratory hospitalizations was similar for infants >3 months of age. These findings are highly supportive

of an effect of maternal vaccination on infant lower respiratory tract disease, even in the absence of laboratory confirmation.

Only four randomized, controlled efficacy trials of influenza vaccine in pregnant women have been conducted – all four have been conducted in Africa and Southeast Asia and funded by the Bill and Melinda Gates Foundation.^{5,6,7,10} This is a remarkable public health effort that has generated much needed data to inform policy and financing decisions in low resource countries. Influenza vaccines given to pregnant women are safe, well-tolerated and immunogenic, and prevent laboratory-confirmed disease in mothers and infants. Importantly, there is a growing body of evidence supporting the impact of these vaccines on other important public health outcomes, including lower respiratory tract disease in infants. In addition to the direct health benefits of influenza vaccine, additional benefits of maternal immunization programs include strengthening antenatal vaccine delivery in preparation for other vaccines (e.g. respiratory syncytial virus) and improving country pandemic preparedness efforts.¹¹ These data are timely and critical to inform upcoming global investment decisions on maternal influenza immunization.

References

1. Ortiz JR, Perut M, Dumolard L. A global review of national influenza immunization policies: Analysis of the 2014 WHO/UNICEF Joint Reporting Form on immunization. *Vaccine*. 2016; 34: 5400-5405.
2. Gessner BD, Brooks WA, Neuzil KM, et al. Vaccines as a tool to estimate the burden of severe influenza in children of low-resourced areas. *Vaccine*. 2013;31(32):3222–28.
3. Feikin DR, Scott JAG, Gessner BD. Use of vaccines as probes to define disease burden. *Lancet* 2014.
4. Cutts FT, Zaman SM, Enwere G, et al. Efficacy of nine-valent pneumococcal conjugate vaccine against pneumonia and invasive pneumococcal disease in The Gambia: randomised, doubleblind, placebo-controlled trial. *Lancet*. 2005;365(9465):1139-1146.
5. Madhi SA, Cutland CL, Kuwanda L, et al. Influenza vaccination of pregnant women and protection of their infants. *N Engl J Med*. 2014;371(10):918-931.
6. Tapia MD, Sow SO, Tamboura B, et al. Maternal immunisation with trivalent inactivated influenza vaccine for prevention of influenza in infants in Mali: a prospective, active-controlled, observer-blind, randomised phase 4 trial. *Lancet Infect Dis*. 2016;16(9):1026-1035.
7. Steinhoff MC, Katz J, Englund JA, et al. Year-round influenza immunisation during pregnancy in Nepal: a phase 4, randomised, placebo-controlled trial. *Lancet Infect Dis*. 2017.
8. Omer SB, Clark DR, Aqil AR, et al. Maternal influenza immunization and prevention of severe clinical pneumonia in young infants: analysis of randomized controlled trials conducted in Nepal, Mali and South Africa. *Pediatr Infect Dis J* 2018 (in press)

9. Nunes MC, Cutland CL, Jones S, et al. Efficacy of maternal influenza vaccination against all-cause lower respiratory tract infection hospitalizations in young infants: Results from a randomized controlled trial. *Clin Infect Dis*. 2017. 65(7):1066-71.
10. Zaman K, Roy E, Arifeen SE, et al. Effectiveness of maternal influenza immunization in mothers and infants. *N Engl J Med*. 2008;359(15):1555-1564.
11. Ortiz JR, Neuzil KM. Influenza immunization of pregnant women in resource-constrained countries: an update for funding and implementation decisions. *Curr Opin Infect Dis* 2017, 30:455–462